

1000, 3000, 5000 metre contours

FIGURE 1. Catch locations of seabirds killed during period 1 October 1996 to 31 December 1997, and returned for autopsy from New Zealand fisheries.

Seabed Mapping
New Ze8Jart9 LIC



(\*u~tnc 1". 1


New Zealand EEZ New Zealand 1000, 3000, 5000 metre contours 0 Diomedea antipodensis

 Diomedea chionoptera
 i Diomedea epomophora Diomedea gibsoni


• Diomedea sanfordi Macronectes halli

O Phoebetria palpebrata


Catch locations during period 1 October 1996 to 31 December 1997, FIGURE 2. of Diomedea, Macronectes and Phoebetria returned for autopsy.




**FIGURE 2.** Catch locations during period 1 October 1996 to 31 December 1997, of Diomedea, Macronectes and Phoebetria returned for autopsy.



**FIGURE 3.** Catch locations during period 1 October 1996 to 31 December 1997, of Thalassarche returned for autopsy.



**FIGURE 4.** Catch locations during period 1 October 1996 to 31 December 1997, of Procellaria petrels returned for autopsy.



**FIGURE 5.** Catch locations during period 1 October 1996 to 31 December 1997, of Puffinus shearwaters returned for autopsy.